Audit 2024

Contact:

Tumorzentrum Hirslanden Zürich Witellikerstrasse 40 8032 Zürich T +41 44 387 96 62 tumorzentrum.hirslanden@hirslanden.ch

open studies

	name of study	target group/ short description
breast cancer	Ascent-03 study	A randomized, open-label, phase 3 study of sacituzumab govitecan versus treatment of
		physician's choice in patients with previously untreated, locally advanced, inoperable or metastatic
		triple-negative breast cancer whose tumors do not express PD-L1 or in patients previously treated
		with anti-PD-(L)1 agents in the early setting whose tumors do express PD-L1

EXPERT- study (ANZ 1601/ BIG 16-02)

HER2CLIMB-05 (Seagen) study

LidERA-study (Roche G042784)

POLAR-study (IBCSG 59/19)

SERENA 4-study (Astra Zeneca)

SASCIA-study (Gilead)

Mallorca-study

PROMs

Ascent-04 study

A randomized, open-label, phase 3 study of sacituzumab govitecan and pembrolizumab versus treatment of physician's choice and pembrolizumab in patients with previously untreated, locally advanced inoperable or metastatic triple-negative breast cancer, whose tumors express PD-L1

EKIB study

Analysis of human breast tumor heterogeneity and immune cell infiltrates and their relevance for patient outcome by mass cytometry.

Cinical trial which presents a unique opportunity to improve personalised use of radiation therapy in patients with early breast cancer according to their individual risks of local recurrence (molecular luminal A-breast cancer)

A study of tucatinib or placebo with trastuzumab and pertuzumab for metastatic HER2+ breast cancer.

A study evaluating the efficacy and safety of adjuvant Giredestrant compared with physician's choice of adjuvant endocrine monotherapy in participants with estrogen receptor-positive, HER2-negative early breast cancer (lidERA Breast Cancer).

Surgical MArker LocaLisation OR Clip and wire Application for targeted axillary dissection in node positive breast cancer patients.

Palbociclib for HR Positive / HER2-negative isolated locoregional recurrence of breast cancer (POLAR).

Patient reported outcomes after oncoplastic breast conserving surgery.

Phase III, prospective, multi-center, randomized, open label, parallel group, study in patients with HER2-negative breast cancer with residual disease after neoadjuvant chemotherapy.

A comparative study of AZD9833 plus Palbociclib versus Anastrozole plus Palbociclib in patients with ER-Positive HER2 negative breast cancer who have not received any systemic treatment for advanced disease (SERENA-4).

	TAXIS-study (SAKK 23/16) VISION-study (SAKK 23/18)	Tailored AXI llary Surgery with or without axillary lymph node dissection followed by radiotherapy in patients with clinically node-positive breast cancer (TAXIS). Vacuum assisted biopsy Immediately before surgery as an Intra- or pre-Operative surrogate for patient response to neoadjuvant chemotherapy for breast cancer (VISION I)
urogenital tumors	PCO study (Prostate Cancer Outcome)	Patient reported outcome measrement of patients with prostate cancer with global uniform patient questionaire.
	Ironman study	The IRONMAN registry collects information about a man's type of prostate cancer, their treatment and what side effects they may be experiencing.
	MRI bladdercancer study	Multiparametric Magnetic Resonance Imaging (MRI) to improve bladder cancer staging and treatment decision-making.
	SAKK 06/ 19 study	Intravesical recombinant BCG followed byperioperative chemo-immunotherapy for patients with MIBC.
gastrointestinal tumors	EDIUM study	Patient-reported outcome study for colon cancer patients: Ergebnisqualität bei D armkrebs: Identifikation von U nterschieden und M assnahmen zur flächendeckenden Qualitätsentwicklung (Quality of Experience in the case of colon cancer: identification of differences und actions to be taken for a comprehnesive quality development)
	ESODATA study	International standardized online database für outcomes of the patients receiving evaluation and/ or treatment für esoohagal diseases of benign and malignant conditions.
	IBDS -CRC study	Image-based ex vivo drug screening and genomics/transcriptomics for patients with colorectal cancer stage II, III or IV
	Patient-derived organoid generation study	Patient-derived organoid generation in pancreatic cancer: a single centre, open-label, single arm feasibility study.
	PREFER-study	Endoscopic management of patients with T1bN0M0 esophageal adenocarcinoma.
	Quality assurance study for duodenal tumors	Quality assurance study of perioperative morbidity and post-surgical follow up of hepato-duodeno- pancreatic surgery.
	Quality assurance study for pancreatic tumors	Quality assurance study of perioperative morbidity and post-surgical follow up of pancreatic surgery.
	ROBOT-2 study	Robot-assisted minimally invasive thoraco-laparoscopic esophagectomy versus minimally invasive esophagectomy for resectable esophageal adenocarcinoma, a randomized controlled trial.
	Tiger study	Distribution of lymph node metastases in esophageal carcinoma.
hematological systemic diseases	Abbvie Valor-Studie	A study to assess change in disease state in adult participants with acute myeloid leukemia (AML) ineligible for intensive chemotherapy in Switzerland and Austria receiving oral Venetoclax tablets (VALOR).
	EBMT-Register	Registry for patients that received a haematopoietic stem cell transplantation or CAR T-cell therapy.
	SAKK EMCL-Registry	The registry of the European Mantle Cell Lymphoma study group - MCL: Disease pattern, treatment algorithms and mechanisms of relapse.

neuro-oncological tumors Gliostar study A study to evaluate safety and efficacy of L19TNF plus lomustine in patients with glioblastoma at

first progression (GLIOSTAR).

Meningiom-study A population-based study on incidence, survival and molecular characteristics of meningioma in

the Canton of Zurich, Switzerland.

PERGOLA-study Pembrolizumab for newly diagnosed glioblastoma: a prospective, open-label, single-arm,

multicenter phase II study.

multi-organ studies AGE-1 Studie Evaluation of forensic methods for age estimation of patient with cancer and a cotrol group.

Roche GLYCART Studie Ex vivo modeling of fresh tumor tissue.